Figure 1. HepG2 cell medium apo B responses after long-term incubation with insulin ± tangeretin





Figure 2. Serum total cholesterol responses in hamsters fed Fructose +0.25% PMF +0.50% PMF +1.00% PMF Control experimental diets Š က շ/լօաա

2/15

Figure 3. HDL cholesterol responses in hamsters fed Fructose +0.25% PMF +0.50% PMF +1.00% PMF Control experimental diets 2 က **၂/၂၀աա**

3/15

Figure 4. Serum triacylglycerol responses in hamsters fed experimental diets



WO 02/087567 PCT/CA02/00662

Figure 5. Serum NEFA responses in hamsters fed experimental diets





Figure 6. Serum insulin responses in hamsters fed experimental diets



WO 02/087567 PCT/CA02/00662

Figure 7. Serum nitrate/nitrite levels in hamsters fed * experimental diets 0.5% PMF 1.0% PMF Control 6 0 1 - 09 80 100 20 140 120






Figure 9. Correlation between serum insulin and nitrate/nitrite levels



10/15


1. In vitro inhibition of alpha-glucosidase by citrus flavonoids and coumarins

Figure 11.



11/15

Figure 12. Effect of experimental diets on serum cholesterol profiles



12/15

Figure 13. Effect of experimental diets on serum triacylglycerols and NEFA




Figure 14. Correlation between serum triacylglycerol and NEFA levels




Figure 15. Glucose tolerance test in hamsters fed experimental diets.

